

λ -HOMOTOPY AND ENGULFABLE HAMILTONIANS

PETER W. SPAETH

ABSTRACT. We apply Lagrangian intersection Floer theory to construct a flat neighborhood of the identity within the Hamiltonian diffeomorphism group of any symplectic manifold with respect to the Hofer and spectral norms.

MSC2000: 53D12; 53D35; 53D40.

1. INTRODUCTION

The purpose of this article is twofold. We show that certain cap moduli spaces entering in the Floer theory of Lagrangian submanifolds are non-empty, a chain level result in the subject. The method of proof follows Chekanov's version [2] of the Floer continuation argument and a generalized definition of chain homotopy, λ -homotopy. The second goal is to show that under the Hofer and spectral norms the Hamiltonian group of a closed symplectic manifold exhibits a local flatness property. This was first observed for the group of Hamiltonian diffeomorphisms of $\mathbb{R}^{2n}$ for the Hofer norm by Bialy-Polterovich [1]. The neighborhood of the identity we construct is likely larger than has been constructed in the literature, notably by McDuff [6] for the the Hofer norm and Y.-G. Oh [11] for both the Hofer and spectral norms. It is defined as follows.

Let (M, ω) be a closed symplectic manifold of dimension $2n$. Following Oh [11] a diffeomorphism ϕ is *engulfable* if it is C^0 -small in the sense that its graph is contained in a Darboux-Weinstein neighborhood of the diagonal and its image under the Darboux chart Φ within the cotangent bundle is the graph of an exact one-form, dS_ϕ , S_ϕ being the (autonomous) generating function for ϕ . Any C^1 -small diffeomorphism is engulfable. Let H^ϕ denote the Hamiltonian function with the property that

$$\Phi(\Delta_{\phi_H^t}) = \text{graph } t \, dS_\phi,$$

where $\Delta_{\phi_H^t}$ is the graph of ϕ inside $(M \times M, -\omega \oplus \omega)$.

This Hamiltonian function is quasi-autonomous [1] and so

$$\int_0^1 \max_{x \in M} H(x, t) - \min_{x \in M} H(x, t) \, dt = \text{osc}(S_\phi) = \max S_\phi - \min S_\phi.$$

The flat neighborhood in question consists of all engulfable Hamiltonians for which the oscillation of the generating function is less than half the minimal energy of a non-constant pseudo-holomorphic sphere in M , what we denote $\sigma(M, \omega)$.

Date: February 2, 2008.

Key words and phrases. Hofer norm, spectral norm, λ -homotopy, Floer theory, thick and thin decomposition.

Let (P, ω) be any tame symplectic manifold and L be a connected, closed and Lagrangian submanifold of P . Also let $\mathcal{U}$ be a Darboux neighborhood of L . Throughout this article we assume all Hamiltonian functions $H : M \times [0, 1] \rightarrow \mathbb{R}$ are normalized. Let $H : P \times [0, 1] \rightarrow \mathbb{R}$ and let $0 < s < 1$. Denote $\sigma(L; P, \omega)$ the minimal area of any non-constant pseudo-holomorphic sphere or disk with boundary on L on P .

Theorem A. *Suppose that the Hamiltonian diffeomorphism $\phi = \phi_H^1$ is non-degenerate in the sense that $\phi_H^1(L)$ meets L transversally and satisfies*

$$(1.1) \quad \|H\| < \frac{\sigma(L; P, \omega)}{2},$$

where $\|H\|$ is Hofer's norm of H . The thin Floer cap product

$$[L] \cap_s^\circ (\cdot) : CF_*(L, sH) \rightarrow CF_*(L, sH)$$

is λ -homotopic to

$$\Phi^- \circ [L] \cap^\mathcal{U} (\cdot) \circ \Phi^+ : CF_*(L, H) \rightarrow CF_*(L, H)$$

where Φ^+ and Φ^- are continuation maps

$$\Phi^+ : CF_*(L, sH) \rightarrow CF_*(L, H) \quad \Phi^- : CF_*(L, H) \rightarrow CF_*(L, sH),$$

and $[L] \cap^\mathcal{U} : CF_*(L, H) \rightarrow CF_*(L, H)$ is the local Floer cap product defined via curves with image in $\mathcal{U}$.

The spectral norm γ on the Hamiltonian group is a Floer theoretic refinement of the Hofer metric and in particular is a lower bound for the Hofer norm. Roughly speaking the spectral norm of a Hamiltonian diffeomorphism ϕ_F is the smallest action difference among critical points of the action functional $\mathcal{A}_F$ of Conley-Zehnder index n and $-n$. Under the isomorphism between Floer homology and quantum (co)homology, such critical points correspondence with the quantum cohomology class 1. To reflect the fact that the spectral metric involves only this cohomology class, Oh introduced a quantity called the homological area $A(\phi; 1)$ which is, again roughly speaking, the minimal energy of a Floer trajectory connecting the maximum and minimum of F . In fact a main result of [11] is that for any non-degenerate Hamiltonian ϕ

$$A(\phi; 1) \leq \gamma(\phi) \leq \|\phi\|_{\text{Hofer}}.$$

In the case of C^1 -small diffeomorphisms the thick and thin decomposition of the Floer moduli space implies $A(\phi; 1) \geq \gamma(\phi)$ and as a result $A(\phi; 1) = \gamma(\phi) = \|\phi\|_{\text{Hofer}} = \text{osc}(S_\phi)$. When the diffeomorphism ϕ is engulfable the thick and thin approach does not directly apply and so we must use Theorem A to prove the following

Theorem B. *Let ϕ be a non-degenerate, engulfable Hamiltonian diffeomorphism and assume that the oscillation of its generating function S_ϕ satisfies*

$$(1.2) \quad \text{osc}(S_\phi) < \frac{\sigma(M, \omega)}{2}.$$

For such a diffeomorphism the following string of equalities holds.

$$(1.3) \quad A(\phi; 1) = \gamma(\phi) = \|\phi\|_{\text{Hofer}} = \text{osc}(S_\phi).$$

Consequently the Lie algebra of the Hamiltonian group can be identified with the normed vector space of normalized C^∞ -functions on M , where the norm is the oscillation. See [6], Proposition 1.8 and [11], Corollary G.

This work formed a portion of my Ph. D. thesis. I wish to thank my advisor Yong-Geun Oh for his guidance and support. Theorem B is motivated by a version of that result announced in the preprint [10]. I also wish to thank Joel Robbin and Augustin Banyaga for many helpful discussions .

2. λ -HOMOTOPY

Chekanov [2] defined an algebraic framework to prove an existence theorem of Lagrangian intersections. We recall the definitions and prove a new algebraic result, Proposition 2.4, needed in the proof of Theorem B.

Let Γ be a free abelian group and $\lambda : \Gamma \rightarrow \mathbb{R}$ be an injective homomorphism. Such a mapping is called a weight function. Denote

$$(2.1) \quad \Gamma^+ = \{g \in \Gamma \mid \lambda(g) > 0\} \quad \Gamma^- = \{g \in \Gamma \mid \lambda(g) < 0\}.$$

Let k be a commutative ring with nonzero unity and $K = k(\Gamma)$ be the group ring over k . The group ring K includes the group Γ as a subgroup.

Now let M be a k -module. The group Γ needn't act on M , but Γ does act on $M \otimes K$ via the second factor $g(v \otimes \bar{g}) := v \otimes g(\bar{g})$. Furthermore $M \otimes K$ admits the decomposition

$$(2.2) \quad M \otimes K = M^+ \oplus M^0 \oplus M^-$$

where $M^+ = \Gamma^+(M \otimes K)$, $M^0 = M$ and $M^- = \Gamma^-(M \otimes K)$.

The above splitting supports two projections

$$p^+ : M \otimes K \rightarrow M^+ \oplus M^0 \text{ and } p^- : M \otimes K \rightarrow M^0 \oplus M^-,$$

$p^+(v \otimes \bar{g}) = v \otimes \sum p^+(\alpha_i)g_i$ where

$$p^+(\alpha_i) = \begin{cases} 0 & \text{if } \lambda(g_i) > 0 \\ \alpha_i & \text{if } \lambda(g_i) \leq 0 \end{cases}$$

and similarly for p^- .

Now assume in addition that (M, ∂) is a differential k -module with $\partial \circ \partial = 0$. Notice that ∂ extends to $M \otimes K$ by setting $\partial(v \otimes \bar{g}) = (\partial v) \otimes \bar{g}$. Also the differential ∂ and the projections p^+, p^- are commuting operators because ∂ acts on M while the projections act only on K in $M \otimes K$.

Definition 2.1 (Chekanov, [2]). *Two K -linear maps $\psi_1, \psi_2 : M \otimes K \rightarrow M \otimes K$ are called λ -homotopic if there exists a K -linear map $h : M \otimes K \rightarrow M \otimes K$ such that*

$$(2.3) \quad p^+(\psi_1 - \psi_2 + h \circ \partial + \partial \circ h)p^- = 0.$$

Notice that if ψ_1, ψ_2 were chain maps in the usual sense that Equation (2.3) holds. The definition implies that ψ_1 and ψ_2 are chain maps from negative to positive weight. The essential algebraic lemma, observed first by Chekanov, from which this section really follows is

Lemma 2.2 (Chekanov, [2]). *Let V be the homology of (M, ∂) . Set $W = V \otimes K$. For any nonzero $r \in W$ there exists $g \in \Gamma$ for which $g \cdot r = q^0 + q^-$, with $q^0 \in V \subset M^0$ nonzero and $q^- \in M^-$.*

Proof. We have $r = v \otimes \bar{g} \in W = V \otimes K$, with $\bar{g} = \sum \alpha_i g_i, \alpha_i \in k, g_i \in \Gamma$. Let $g = g_j \in \Gamma$ be such that $\lambda(g) = -\max_{\alpha_i \neq 0} \lambda(g_i)$. The quantity $\max_{\alpha_i \neq 0} \lambda(g_i)$ is called the *valuation* of g . Then by the action of Γ on K $g(r) = v \otimes \sum_{g_n g_j = g_i} \alpha_n g_i$. Finally if $\alpha_n \neq 0$ then $\lambda(g_i) = \lambda(g_n) + \lambda(g_j) \leq -\lambda(g_j) + \lambda(g_j) = 0$ and hence $\lambda(g_i) \leq 0$. $\square$

As an example of what can be proven using this lemma we state the following

Lemma 2.3 (Lemma 5, [2]). *Let λ, Γ, k, K and (M, ∂) be as above. Let in addition N be a K -module and the maps $\Phi^+ : M \otimes K \rightarrow N, \Phi^- : N \rightarrow M \otimes K$ be K -linear. If $\Phi^- \circ \Phi^+$ is λ -homotopic to the identity, then $\text{rank}_K(N) \geq \text{rank}_k H(M, \partial)$, where $H(M, \partial)$ is the homology of the complex (M, ∂) .* $\square$

What follows below is the main algebraic result of the current paper. An isomorphism on homology can not be λ -homotopic to the zero map provided the isomorphism commutes with the projection p^+ .

Proposition 2.4. *Suppose (M, ∂) is a differential k -module with non-trivial homology. Suppose the linear maps $\psi_1, \psi_2 : M \otimes K \rightarrow M \otimes K$ are λ -homotopic. If ψ_1 descends to an isomorphism $\psi_1 : H(M, \partial) \xrightarrow{\cong} H(M, \partial)$ and the induced map commutes with p^+ on homology (i.e. up to a boundary on the chain level) then $\psi_2 \neq 0$.*

Proof. Choose a submodule of ∂ -cycles $V \subset M$ representing the homology $H(M, \partial)$. Let $W = V \otimes K$. Note that W is not trivial. By Lemma (2.2) for any nonzero $r \in W$ there exists $g \in \Gamma$ for which $g \cdot r = q^0 + q^-$, with $q^0 \in V \subset M^0$ nonzero and $q^- \in M^-$.

To prove the proposition suppose to the contrary that $\psi_2 = 0$. Applying equation (2.3) to $q^0 + q^-$ we see

$$\begin{aligned} 0 &= p^+ (\psi_1 - \psi_2 + h \circ \partial + \partial \circ h) p^- (q^0 + q^-) \\ &= p^+ (\psi_1 - \psi_2 + h \circ \partial + \partial \circ h) (q^0 + q^-) \\ &= p^+ (\psi_1 (q^0 + q^-)) + p^+ (h \circ \partial + \partial \circ h) (q^0 + q^-) \\ &= p^+ (\psi_1 (q^0 + q^-)) + \partial (p^+ h (q^0 + q^-)) \end{aligned}$$

Here we have used the linearity of all the maps, that $q^0 + q^-$ is a cycle and that the projection p^+ commutes with the boundary. By assumption p^+ and ψ_1 commute up to a boundary, say ∂c and so continuing

$$\begin{aligned} 0 &= \psi_1 (q^0) + \partial c + \partial (p^+ h (q^0 + q^-)) \\ &= \psi_1 (q^0) + \partial (c + p^+ h (q^0 + q^-)) \end{aligned}$$

which implies that $\psi_1 (q^0)$ is a boundary; hence ψ_1 can not be an isomorphism, a contradiction. $\square$

Remark 2.5. Because the valuation of any element in the downward Novikov ring is finite, Lemma (2.2) continues to hold and so λ -homotopy extends nicely to this coefficient ring.

3. THE PROOF OF THEOREM A

Let L be a closed, connected n -dimensional Lagrangian submanifold of the tame symplectic (P, ω) . Let $\mathcal{U}$ be a Darboux-Weinstein neighborhood of L . We employ the version of Floer homology of the action functional from [7], [5]. Fix a

reference path l_0 on L . Given a (normalized) time-dependent Hamiltonian function $H : P \times [0, 1] \rightarrow \mathbb{R}$ the corresponding action functional $\mathcal{A}_H$ is real valued on the Novikov covering $\tilde{\Omega}_0(L, l_0; P, \omega) = \tilde{\Omega}_0 \xrightarrow{\pi} \Omega_0$ of the space Ω_0 of contractible paths beginning and ending on L . $\tilde{\Omega}_0$ consists of equivalence classes $[l, w]$ where the half disk w provides the homotopy to l_0 and two pairs (l_1, w_1) and (l_2, w_2) are equivalent if $l_1 = l_2$ and the Maslov index μ_L and symplectic action I_ω homomorphisms vanish on the glued disk $w_1 \# \overline{w_2}$.




FIGURE 3.1. Half-disk w with boundary on L

The choice of sign is

$$\mathcal{A}_H([l, w]) := - \int w^* \omega - \int_0^1 H(l(t), t) dt.$$

Denote $\Gamma = \Gamma(L, l_0; P, \omega)$ the automorphism group of this cover, a free abelian group. Γ acts by gluing a disk: $g \cdot [l, w] = [l, w \# g]$. The symplectic action provides the weight homomorphism

$$\lambda(g) := - \int g^* \omega.$$

Let $\sigma(L; P, \omega)$ be the minimal area of a pseudo-holomorphic sphere or disk with boundary on L in P and assume for the remainder of this section that the Hofer norm of H satisfies $\|H\| < \sigma/2$. Let $0 < s \ll 1$ and choose $0 < \delta$ to satisfy $\|H\| < \delta/2 < \sigma/2$.

For the parametrized Hamiltonian sH denote the critical points of the action functional $\mathcal{A}_{sH}$

$$\text{Crit}(\mathcal{A}_{sH}) = CF(L, sH) = CF(s)$$

and the vector space of Floer chains

$$CF_*(s) = CF_*(L, sH; \mathbb{Z}_2) = \text{span}_{\mathbb{Z}_2}(CF(s)).$$

We will make a number of assumptions on the size of the parameter s in the course of the proof of Theorem A, the first of which is

$$(3.1) \quad \phi_{sH}^t(L) \subset \mathcal{U} \quad (\forall) \ t \in [0, 1].$$

Two elements of $CF(s)$ are equivalent [2] if they belong to the same connected component of the set

$$\pi^{-1}\{l \in \Omega_0 : l([0, 1]) \subset \mathcal{U}\} \subset \tilde{\Omega}_0.$$

The first assumption on the parameter s guarantees that the set of equivalence classes is not empty. Fixing any equivalence class $CF^\circ(s)$ yields the decomposition $CF_*(s) = CF_*^\circ(s) \otimes K$, where $CF_*^\circ(s) = \text{span}_{\mathbb{Z}_2} CF^\circ(s)$ and $K = \mathbb{Z}_2(\Gamma)$ denotes the group ring. The action difference of any two elements in $CF^\circ(s)$ can be made arbitrarily small by choosing s sufficiently small. This is the second assumption we make on s :

$$(3.2) \quad [x, \tilde{x}], [z, \tilde{z}] \in CF^\circ(s) \implies \mathcal{A}_{sH}([x, \tilde{x}]) - \mathcal{A}_{sH}([z, \tilde{z}]) < \delta/2$$

The sets of positive and negative weight critical points are denoted

$$CF^+(L, sH) = \Gamma^+(CF^\circ) \text{ and } CF^-(L, sH) = \Gamma^-(CF^\circ).$$

Finally we fix our notation for the Floer moduli spaces. Let J be a (possibly time-dependent) almost complex structure compatible to ω . The Floer differential is defined by finite energy solutions of the following boundary problem

$$(3.3) \quad \begin{cases} \frac{\partial u}{\partial \tau} + J(u, t) \left(\frac{\partial u}{\partial t} - X_{sH}(u, t) \right) = 0 \\ u : \mathbb{R} \times [0, 1] \rightarrow P \\ u(\tau, 0), u(\tau, 1) \subset L \end{cases}$$

The energy of a solution u is

$$E_J(u) := \int_{-\infty}^{+\infty} \left| \frac{\partial u}{\partial \tau} \right|_J^2 d\tau = \int_{-\infty}^{+\infty} \int_0^1 \omega \left(\frac{\partial u}{\partial \tau}, J \frac{\partial u}{\partial \tau} \right) dt d\tau.$$

The full Floer moduli space of finite energy solutions $\mathcal{M}(L, sH) = \mathcal{M}(s)$ ranges over all critical points of $\mathcal{A}_{sH}$. When asymptotic limits are specified, we write $\mathcal{M}([x, \tilde{x}], [z, \tilde{z}], s)$. For generic s this is a smooth manifold of dimension $\mu_L([x, \tilde{x}]) - \mu_L([z, \tilde{z}])$, when it is not empty.

The length of a Floer trajectory $u \in \mathcal{M}([x, \tilde{x}], [z, \tilde{z}], s)$ is defined to be the action difference at its ends

$$(3.4) \quad l(u) := \mathcal{A}_{sH}([x, \tilde{x}]) - \mathcal{A}_{sH}([z, \tilde{z}]).$$

We say that u is *somewhat thin* if $l(u) \leq ||H|| + \delta$, *thin* if $l(u) \leq \delta$ and local if $\text{image}(u) \subset \mathcal{U}$. Curves which exit the Darboux-Weinstein neighborhood are called *thick*. A consequence of Gromov's compactness theorem is Oh's thick and thin decomposition theorem that implies when s is sufficiently small and J is C^1 -close to a time independent compatible almost complex structure every somewhat thin trajectory is thin and every (somewhat) thin trajectory is local. Our fourth assumption on the size of the parameter s is that it is small enough to ensure the thick and thin decomposition. See [7] and [2]. We denote the collection of thin trajectories $\mathcal{M}^\circ(L, sH)$ and local trajectories $\mathcal{M}^u(L, sH)$. After dividing by the time-shift, the thin Floer differential ∂° is defined by $\mathcal{M}^\circ/\mathbb{R} := \widehat{\mathcal{M}}^\circ(L, sH)$

$$\partial^\circ([x, \tilde{x}]) = \sum_{\mu_L([x, \tilde{x}]) - \mu_L([z, \tilde{z}]) = 1} \#_{\mathbb{Z}_2} \widehat{\mathcal{M}}^\circ([x, \tilde{x}], [z, \tilde{z}], s)[z, \tilde{z}]$$

The thin differential satisfies $\partial^\circ \circ \partial^\circ = 0$ and in fact represents the singular homology of L

$$(3.5) \quad H(CF_*^\circ(s), \partial^\circ) := HF_*^\circ(L, sH; \mathbb{Z}_2) \cong H_*(L; \mathbb{Z}_2).$$

The proof of this isomorphism is carried out in Section 4 of [7].

The thin Floer cap product $[L] \cap_s^\circ (\cdot) : CF_*^\circ(s) \rightarrow CF_*^\circ(s)$ is defined by counting elements $u \in \mathcal{M}^\circ(L, sH)$ passing through a generic point $p \in L$.

$$\mathcal{M}^\circ(L, sH; p) = \{u \in \mathcal{M}^\circ \mid u(0, 0) = p\}$$

$$[L] \cap_s^\circ ([x, \tilde{x}]) = \sum_{\mu_L([x, \tilde{x}]) - \mu_L([z, \tilde{z}]) = n} \#_{\mathbb{Z}_2} \mathcal{M}^\circ([x, \tilde{x}], [z, \tilde{z}], s; p)[y, \tilde{y}]$$

The collection of such maps is denoted $\mathcal{M}^\circ(L, sH; p)$ (or $\mathcal{M}^\circ([x, \tilde{x}], [z, \tilde{z}], s; p)$ or $\mathcal{M}^\mathcal{U}(L, sH; p)$, etc.). Because of the pointed condition $u(0, 0) = p$, the dimension is reduced by n , the dimension of L . Again the thick and thin decomposition holds.

Proposition 3.1 (Proposition 4.6, [7]). *Fix a time-independent compatible almost complex structure J_0 on P . There exists $s' > 0$ with the property that for any $0 < s < s'$ if $\|J - J_0\|_{C^0(P \times [0, 1])} < s$ then any somewhat thin cap trajectory is thin and any (somewhat) thin cap trajectory is local.*

The Proposition implies that the moduli space $\mathcal{M}^\circ(L, sH; p)$ is isolated within $\mathcal{M}(L, sH; p)$. Passing to the cotangent bundle, the work of [3], [9] and [4] implies $[L] \cap_s^\circ (\cdot)$ descends to an isomorphism on $HF_*^\circ(L, sH; \mathbb{Z}_2)$. (In particular A. Floer originally proved the injectivity of the cap action.) The next assumption on s is that it is small enough to ensure

$$(3.6) \quad [L] \cap_s^\circ (\cdot) : HF_*^\circ(L, sH; \mathbb{Z}_2) \xrightarrow{\cong} HF_{*-n}^\circ(L, sH; \mathbb{Z}_2).$$

Setting the parameter $s = 1$, the local cap product $[L] \cap^\mathcal{U} (\cdot)$ is defined by the local curves in $\mathcal{M}^\mathcal{U}(L, H; p)$.

$$[L] \cap^\mathcal{U} ([y, \tilde{y}]) = \sum_{\mu_L([y, \tilde{y}]) - \mu_L([y', \tilde{y}']) = n} \#_{\mathbb{Z}_2} \mathcal{M}^\mathcal{U}([y, \tilde{y}], [y', \tilde{y}'], 1; p)[y', \tilde{y}']$$

Within the Darboux neighborhood of L bubbling-off can not occur and so $[L] \cap^\mathcal{U} (\cdot)$ is defined; however, this map need not agree with the singular cap product. We have thus far fallen short in proving that the moduli space of local curves is isolated among all curves. This fact is what motivated the use of λ -homotopy. For the same reason, when $s = 1$ the local Floer homology might not be defined. In particular $\partial^\mathcal{U}$ need not square to zero.

The continuation maps Φ^+ and Φ^- are constructed by introducing monotone continuation functions ρ^+ and ρ^- of τ and a non-autonomous Floer equation. We sketch the map Φ^+ , referring the reader to [2] for the full details.

Let $b_+(H) = \int_0^1 \max_x H(x, t) dt$ and $b_-(H) = - \int_0^1 \min_x H(x, t) dt$. Let ρ^+ be a smooth increasing function of τ so that $\rho^+(\tau) = 1$ for $\tau \gg 1$ and $\rho^+(\tau) = s$ for $\tau \ll -1$. Consider the continuation equation

$$(3.7) \quad \begin{cases} \frac{\partial u}{\partial \tau} + J^{\rho^+}(u, t) \left(\frac{\partial u}{\partial t} - X_{\rho^+ H}(u, t) \right) = 0 \\ u : \mathbb{R} \times [0, 1] \rightarrow P \\ u(\tau, 0), u(\tau, 1) \subset L \end{cases}$$

and define

$$\mathcal{M}_{\rho+} = \{u : \mathbb{R} \times [0, 1] \rightarrow P \mid E_J(u) < +\infty \text{ \& (3.7)}\}$$

The length of a continuation trajectory can be defined as before (and could be negative) $u^+ \in \mathcal{M}_{\rho+}$

$$l(u^+) = \mathcal{A}_{sH}([x, \tilde{x}]) - \mathcal{A}_H([y, \tilde{y}])$$

where $u^+(-\infty) = [x, \tilde{x}] \in CF(s)$ and $u^+(+\infty) = [y, \tilde{y}] \in CF(1)$. The trajectory u^+ is called thin if $l(u^+) \leq (1-s) \cdot b_+(H) + \delta$ and the collection of all thin trajectories we denote $\mathcal{M}_{\rho+}^\circ$.

Lemma 3.2 (Lemma 9, [2]). *If $u^+ \in \mathcal{M}_{\rho+}$, then $l(u^+) \geq -(1-s) \cdot b_-(H)$. If $u^- \in \mathcal{M}_{\rho-}$, then $l(u^-) \geq -(1-s) \cdot b_+(H)$. For the short continuation trajectories $E(u^\pm) \leq \|H\| + \delta$.*

Consequently the continuation map $\Phi^+ : CF(s) \rightarrow CF(1)$ defined by

$$\Phi^+([x, \tilde{x}]) = \sum \#_{\mathbb{Z}_2} \mathcal{M}_{\rho+}^\circ([x, \tilde{x}], yy)[y, \tilde{y}]$$

is finite. Notice that a thick and thin decomposition is not needed to make this definition.

The important point is that these linear maps provide a bridge from the Floer complexes corresponding to $0 < s < 1$ and $s = 1$.

$$(3.8) \quad \Phi^+ : CF_*(L, sH) \rightarrow CF_*(L, H) \text{ and } \Phi^- : CF_*(L, H) \rightarrow CF_*(L, sH)$$

The main result in [2] is that when s is sufficiently small and $\|H\| < \sigma$, there exists a λ -homotopy h from the identity to the composition $\Phi^- \circ \Phi^+$. As a consequence the displacement energy of any Lagrangian submanifold is larger than σ . Our λ -homotopy is defined similarly, where we must also take into consideration the marked point $p \in L$. We now provide the details.

Let $R \in [s, +\infty)$ and $\mu_R : \mathbb{R} \rightarrow (s, 1]$ be a smooth function with all the following properties. Firstly $0 \leq \mu'_R(\tau)$ for $\tau \leq 0$ and $\mu'_R(\tau) \leq 0$ for $\tau \geq 0$ and for $R \geq 1$

$$(3.9) \quad \mu_R(\tau) = \begin{cases} 1 & -R \leq \tau \leq R \\ s & |\tau| \geq R+1 \end{cases}$$

while when $s \leq R \leq 1$, $\mu_R = R \cdot \mu_1$. The λ -homotopy is defined via the the finite energy solutions of the non autonomous Floer equation

$$(3.10) \quad \begin{cases} \frac{\partial u}{\partial \tau} + J(u, t) \left(\frac{\partial u}{\partial t} - X_{\mu_R(\tau)}(u, t) \right) = 0 \\ u(\tau, 0), u(\tau, 1) \subset L \end{cases}$$

The collection of all finite energy solutions is denoted $\mathcal{M}_{\mu_R} = \mathcal{M}_{\mu_R}(L, sH)$. Then define the parametrized moduli space $\mathcal{M}_\mu = \{(R, v) : v \in \mathcal{M}_{\mu_R}\}$. By the parameterized index theorem the local dimension of $\mathcal{M}_\mu$ is one greater than the difference of Maslov indices of the asymptotic limits and so defines a map of degree +1. The formula for the pointed λ -homotopy $h^p : CF(s) \rightarrow CF(s)$ on generators is

$$(3.11) \quad h^p([x, \tilde{x}]) = \sum_{[y, \tilde{y}] \in CF^\circ \cup CF^+} \#_{\mathbb{Z}_2} \{\mathcal{M}_\mu([x, \tilde{x}], [y, \tilde{y}]; p)\} [y, \tilde{y}]$$

where $[x, \tilde{x}] \in CF^- \cup CF^\circ$. Because the energy is controlled by the Hofer norm, bubbling off does not occur and so the sum is finite. The induced map is linear over K .

We now verify the λ-homotopy equation.

$$(3.12) \quad p^+([L] \cap_s^\circ (\cdot) - \Phi^- \circ [L] \cap^{\mathcal{U}} (\cdot) \circ \Phi^+ + h^p \circ \partial^\circ + \partial^\circ \circ h^p)p^- = 0$$

for all $[x, \tilde{x}] \in CF^\circ(s)$. We write the left hand side of Equation (3.12) as

$$\sum \#_{\mathbb{Z}_2} \mathcal{S}([x, \tilde{x}], [z, \tilde{z}])[z, \tilde{z}]$$

where $\mathcal{S}([x, \tilde{x}], [z, \tilde{z}])$ is the union of the following four sets. For the remainder of this section each of $[x, \tilde{x}] \in CF^\circ(s) \cup CF^-(s)$, $[z, \tilde{z}] \in CF^+(s) \cup CF^\circ(s)$ and $[y, \tilde{y}], [y', \tilde{y}'] \in CF(1)$ represent generic critical points of the respective action functionals.

- (1) A thin cap trajectory $u \in \mathcal{M}^\circ([x, \tilde{x}], [z, \tilde{z}], s; p)$
- (2) A triple (u^-, v, u^+) with $u^+ \in \mathcal{M}_{\rho^+}^\circ([x, \tilde{x}], [y, \tilde{y}])$, $v \in \mathcal{M}^{\mathcal{U}}([y, \tilde{y}], [y', \tilde{y}']; p)$ and $u^- \in \mathcal{M}_{\rho^-}^\circ([y', \tilde{y}'], [z, \tilde{z}])$
- (3) A thin unparametrized gradient trajectory $u \in \widehat{\mathcal{M}}^\circ([x, \tilde{x}], [z, \tilde{z}], s)$ and element (R, v) with $v \in \mathcal{M}_{\mu_R}([x, \tilde{x}], [z, \tilde{z}]; p)$
- (4) An element (R, v) with $v \in \mathcal{M}_{\mu_R}([x, \tilde{x}], [z, \tilde{z}]; p)$ and a thin unparametrized gradient trajectory $u \in \widehat{\mathcal{M}}^\circ([x, \tilde{x}], [z, \tilde{z}], s)$

By the Floer gluing theorem each term above lies at the end of the one-dimensional portion $\mathcal{Q}$ of $\mathcal{S}$. In fact $\mathcal{Q}$ is compact and the correspondence is one-one. The compactness of $\mathcal{Q}$ follows from the uniform energy estimate which is a consequence of the assumption $\|H\|_{Hofer} < \sigma/2$ (See Lemma 10, [2] and Lemma 2.2, [8].).


On the other hand by the Gromov-Floer compactness theorem every end of $\mathcal{Q}$ may be compactified by one of the following

- (1) A cap trajectory $u \in \mathcal{M}^\circ([x, \tilde{x}], [z, \tilde{z}], s; p)$
- (2) A triple (u^-, v, u^+) with $u^+ \in \mathcal{M}_{\rho^+}([x, \tilde{x}], [y, \tilde{y}])$, $v \in \mathcal{M}^{\mathcal{U}}([y, \tilde{y}], [y', \tilde{y}']; p)$ and $u^- \in \mathcal{M}_{\rho^-}([y', \tilde{y}'], [z, \tilde{z}])$
- (3) An unparametrized gradient trajectory $u \in \widehat{\mathcal{M}}([x, \tilde{x}], [z, \tilde{z}], s)$ and element (R, v) with $v \in \mathcal{M}_{\mu_R}([x, \tilde{x}], [z, \tilde{z}]; p)$
- (4) An element (R, v) with $v \in \mathcal{M}_{\mu_R}([x, \tilde{x}], [z, \tilde{z}]; p)$ and an unparametrized gradient trajectory $u \in \widehat{\mathcal{M}}([x, \tilde{x}], [z, \tilde{z}], s)$

We must show that in each of the above cases the relevant trajectories are thin. In fact this follows from the conjugation by p^+ and p^- and that $\|H\| < \sigma/2$. Moreover that the limit trajectories are thin in cases (3) and (4) already follows from Chekanov's original argument. Therefore we focus on (1) and (2).

Let us begin with (1). The asymptotic limits assure us that $[x, \tilde{x}] = g_x \cdot [x', \tilde{x}']$ and that $[z, \tilde{z}] = g_z \cdot [z', \tilde{z}']$ with

$$-\int (g_z \# \tilde{z})^* \omega \leq 0 \text{ and } \int (g_x \# \tilde{x})^* \omega \leq 0.$$

FIGURE 3.2. u^+, u, u^- and their respective asymptotic limits

Hence the length of u satisfies

$$\begin{aligned}
l(u) &= \mathcal{A}_{sH}([z, \tilde{z}]) - \mathcal{A}_{sH}([x, \tilde{x}]) \\
&= - \int \tilde{z}^* \omega - s \int H(z, t) dt + \int \tilde{x}^* \omega + s \int H(x, t) dt \\
&= - \int (g_z \# \tilde{z}')^* \omega - s \int H(z, t) dt + \int (g_x \# \tilde{x}')^* \omega + s \int H(x, t) dt \\
&\leq s \left(\int H(x, t) - H(z, t) dt \right) \\
&\leq s \left(\int \max_x H(x, t) - \min_x H(x, t) dt \right) \\
&= s \|H\| \\
&\leq \delta
\end{aligned}$$

which makes it thin.

Figure 3.2 illustrates the situation of case (2).

To complete the identification we must prove that each u^+ and u^- is thin. In other words we must prove $l(u^+) \leq (1-s) \cdot b_+(H) + \delta$ and $l(u^-) \leq (1-s) \cdot b_-(H) + \delta$. Toward that end, consider the quantity $l(u^+) = l(u^-)$. By definition of length

$$\begin{aligned}
l(u^+) + l(u^-) &= \mathcal{A}_{sH}([z, \tilde{z}]) - \mathcal{A}_H([y, \tilde{y}]) + \mathcal{A}_H([y', \tilde{y}']) - \mathcal{A}_{sH}([z, \tilde{z}]) \\
&= \int g_x^* \omega - s \int H(x, t) dt + \left(\mathcal{A}_H([y', \tilde{y}']) - \mathcal{A}_H([y, \tilde{y}]) \right) \\
&\quad - \int g_z^* \omega + s \int H(z, t) dt \\
&\leq s \left(\int H(z, t) dt - \int H(x, t) dt \right) + E(u) \\
(3.13) \quad &\leq \delta/2 + \delta/2.
\end{aligned}$$

For the last line $E(u) < \|H\| \leq \delta/2$ and from this it now follows that u^+ is thin: Recall from Lemma 3.2 that $l(u^-) \geq -(1-s) \cdot b_+(H)$ (resp. $l(u^+) \geq -(1-s) \cdot b_-(H)$) and so if u^+ were not thin then

$$\delta = -(1-s) \cdot b_+(H) + \delta + (1-s) \cdot b_+(H) < l(u^+) + l(u^-) \leq \delta$$

a contradiction. Repeating the argument but interchanging u^+ and u^- proves that u^- is also thin. In short the terms in Equation (3.12) are in one-one correspondence with the ends of the compact one-dimensions manifold $\mathcal{Q}$. This implies h^p is indeed a λ -homotopy.

4. THE PROOF OF THEOREM B

Theorem B is a consequence of Theorem A and Proposition 2.4. Take the symplectic manifold $(M \times M, -\omega \oplus \omega)$ and the diagonal, Δ , a Lagrangian submanifold of dimension $2n = \dim(M)$. In this case $\sigma(\Delta; M \times M, -\omega \oplus \omega) = \sigma(M; \omega)$. Let $\bar{J}$ be an almost complex structure on $M \times M$ compatible to $-\omega \oplus \omega$ and of the form $-J_{2t-1} \oplus J_{2(1-t)}$ in $\mathcal{U}$.

Throughout this section consider the Hamiltonian deformation of the diagonal corresponding to the isotopy $t \mapsto \text{graph } tdS_\phi$, and assume that the Hamiltonian function H satisfies $\|H\| < \sigma(M, \omega)/2$. For any non-degenerate Hamiltonian diffeomorphism we have $A(\phi; 1) \leq \gamma(\phi) \leq \|\phi\|_{\text{Hofer}} \leq \|H\| = \text{osc}(S_\phi)$. Examination of Oh's proof in the C^1 -small case makes clear that the essential step of the proof is to produce a Floer trajectory connecting the maximum and minimum of the quasi-autonomous Hamiltonian H . The trajectory we find is in fact local. Once this is established we will then have $\text{osc}(S_\phi) \leq A(\phi; 1)$, completing the proof.

When s is sufficiently small the thin part of the Floer homology satisfies

$$HF_{n-j}^\circ(\Delta, sH, \mathbb{Z}_2) \cong H_j(\Delta, \mathbb{Z}_2).$$

Furthermore the thin part of the Floer cap action $[\Delta] \cap_s^\circ(\cdot)$ agrees with the standard cap action of $[\Delta]$, an isomorphism. Let $j = 2n$ and 0 and consider the following diagram

$$\begin{array}{ccc} HF_{-n}^\circ(\Delta, sH; \mathbb{Z}_2) & \xrightarrow{\cong} & H_{2n}(\Delta; \mathbb{Z}_2) \\ [\Delta] \cap_s^\circ(\cdot) \downarrow \cong & & \cong \downarrow [\Delta] \cap(\cdot) \\ HF_n^\circ(\Delta, sH; \mathbb{Z}_2) & \xrightarrow{\cong} & H_0(\Delta; \mathbb{Z}_2) \end{array}$$

Because Δ is orientable $H_0(\Delta; \mathbb{Z}_2) \cong \mathbb{Z}_2$ and $H_{2n}(\Delta; \mathbb{Z}_2) \cong \mathbb{Z}_2$ and so $[\Delta] \cap_s^\circ(\cdot) : \mathbb{Z}_2 \xrightarrow{\cong} \mathbb{Z}_2$ must be the identity. Hence this map commutes with the projection p^+ and Proposition 2.4 implies in combination with Theorem A that the composition $\Phi^- \circ [\Delta] \cap^\mathcal{U}(\cdot) \circ \Phi^+$ is non-zero. Because all three maps are linear this means that the local Floer cap product $[\Delta] \cap^\mathcal{U}(\cdot)$ is non-zero and therefore the moduli space $\mathcal{M}^\mathcal{U}([x^+, \widehat{x^+}], [x^-, \widehat{x^-}]; p)$ must contain an odd number of elements, where $x^\pm$ are respectively the maximum and minimum points of H and $\widehat{x^\pm}$ are constant bounding

disks. In particular it is not empty. Let us denote this element v which satisfies

$$\begin{cases} \frac{\partial v}{\partial \tau} + \overline{J}(v, t) \left(\frac{\partial v}{\partial t} - X_H(v) \right) = 0 \\ v : \mathbb{R} \times [0, 1] \rightarrow M \times M \quad v(\tau, 0), v(\tau, 1) \subseteq \Delta \\ \lim_{\tau \rightarrow \pm\infty} v(\tau, \cdot) = x^\pm \\ v(0, 0) = p \in \Delta \quad \text{image}(v) \subset \mathcal{U} \end{cases}$$

Integration by parts yields $E_{\overline{J}}(v) = \text{osc}(S_\phi)$. We glue the trajectory v into one into M , $u : \mathbb{R} \times S^1 \rightarrow M$, as follows

$$u(\tau, t) = \begin{cases} v(2\tau, 2t - 1) & 0 \leq t \leq 1/2 \\ v(2\tau, 2(1 - t)) & 1/2 \leq t \leq 1 \end{cases}$$

This u is smooth by elliptic regularity and satisfies

$$\begin{cases} \frac{\partial u}{\partial \tau} + J(v, t) \left(\frac{\partial u}{\partial t} - X_H(u) \right) = 0 \\ u(\tau, t) = u(\tau, t + 1) \\ \lim_{\tau \rightarrow \pm\infty} u(\tau, \cdot) = x^\pm(\cdot) \\ u(0, 0) = p. \end{cases}$$

A straightforward chain rule calculation shows that $E_J(u) = E_{\overline{J}}(v) = \text{osc}(S_\phi)$. The final argument in the proof of Theorem 7.1, [11] now applies (See Section 9.4, [11].) and we conclude $\text{osc}(S_\phi) \leq A(\phi, 1)$. This finishes the proof of Theorem B. $\square$

It can be very difficult to compute the constant σ . For the two-sphere with total surface area equal to 4π then in fact $\sigma(S^2 \times S^2, -\omega \oplus \omega)/2 = 2\pi$. The Hamiltonian function which generates the half-turn rotation about a diameter has Hofer length 2π . In this sense the constant $\sigma/2$ seems to be precise. This is plausible as rotation by more than half the way around a diameter is not Hofer (nor spectral) length minimizing: one can generate the same diffeomorphism by rotation in the opposite direction. However it is not yet known if this situation is engulfable. In some cases the Darboux neighborhood of the diagonal can be very large. This will be studied in subsequent work.

REFERENCES

1. M. Bialy and L. Polterovich, *Geodesics of Hofer's metric on the group of Hamiltonian diffeomorphisms*, Duke Math. J. **76** (1994), 273–292.
2. Yuri V. Chekanov, *Lagrangian intersections, symplectic energy, and areas of holomorphic curves*, Duke Math. J. (1998), 213–226.
3. A. Floer, *Morse theory for Lagrangian intersections*, J. Diff. Geom. **28** (1988), 513–547.
4. K. Fukaya and Y.-G. Oh, *Zero-loop open strings and in the cotangent bundle and Morse homotopy*, Asian J. Math. (1997), no. 1, 90–180.
5. K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian Intersection Floer Theory - Anomaly and Obstruction*, manuscript, 2006.
6. D. McDuff, *Geometric variants of the Hofer norm*, J. Symplectic Geom. **1** (2002), 197–252.
7. Y.-G. Oh, *Floer cohomology, spectral sequences, and the Maslov class of Lagrangian embeddings*, Inter. Math. Res. Notices (1996), no. 6, 305–346.
8. ———, *Gromov-Floer theory and disjunction energy of compact Lagrangian embeddings*, Math. Res. Letters **4** (1997), 895–905.
9. ———, *Symplectic topology as the geometry of the action functional, II – pants product and cohomological invariants*, Comm. Anal. Geom. **7** (1999), no. 1, 1–55.
10. ———, *Spectral invariants and the geometry of Hamiltonian diffeomorphisms*, (2004), preprint, arXiv: math.SG/0403083 v1.

11. ———, *Spectral invariants, analysis of the Floer moduli space, and geometry of the Hamiltonian diffeomorphism group*, Duke Math. J. **130** (2005), 199–295.

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, USA

E-mail address: `spaeth@math.psu.edu`